

12

EUROPEAN PATENT APPLICATION

21 Application number: 84303304.4

51 Int. Cl.⁴: A 61 K 9/10

A 61 K 47/00, A 61 K 31/557

22 Date of filing: 16.05.84

30 Priority: 20.05.83 JP 88760/83

43 Date of publication of application:
23.01.85 Bulletin 85/4

84 Designated Contracting States:
BE DE FR GB NL SE

71 Applicant: TAISHO PHARMACEUTICAL CO. LTD
24-1 Takata 3-chome Toshima-ku
Tokyo 171(JP)

71 Applicant: THE GREEN CROSS CORPORATION
15-1, Imabashi-1-chome Higashi-ku Osaka-shi
Osaka 541(JP)

72 Inventor: Mitzushima, Yutaka c/o SEI MARIANNA
IKADAIGAKU
2095 Sugao Takatsu-ku
Kawasaki-shi(JP)

72 Inventor: Aihara, Hironaka
1514-40, Kitanakamaru
Kitamoto-shi(JP)

72 Inventor: Otomo, Susumu
785-341, Kamioidezuka
Konosu-shi(JP)

72 Inventor: Yokoyama, Kazumasa
7 Terauchi-2-chome
Toyonaka-shi(JP)


72 Inventor: Okamoto, Hiroyuki
9-18 Asagiricho-2-chome
Akashi-shi(JP)

72 Inventor: Suyama, Tadakazu
3-7 Matsugaoka-4-chome Tanabecho
Tsuzuki-gun Kyoto(JP)

74 Representative: Harrison, David Christopher et al,
MEWBURN ELLIS & CO 2/3 Cursitor Street
London EC4A 1BQ(GB)

54 Fat emulsion containing prostaglandin.

57 A fat emulsion contains a prostaglandin E₁ alkyl ester represented by the general formula



wherein R denotes an alkyl group having 1 to 30 carbon atoms.

The fat emulsion can be administered intravenously, has a long half-life of its effective ingredient, prostaglandin E₁ alkyl ester, in the living body as well as a focus selectivity.

EP 0 132 027 A1

FAT EMULSION CONTAINING PROSTAGLANDIN

1 This invention relates to a fat emulsion contain-
ing prostaglandin. More particularly, it relates to a fat
emulsion containing prostaglandin E₁ alkyl ester.

 Prostaglandins (hereinafter referred to briefly
5 as PGs) have diversified physiological actions including
vasodilative action, improvement of peripheral blood
circulation, hypotensive action, antilipolysis and
natriuresis, and hence their application to pharmaceuticals
have been investigated for some time past.

10 However, when the potentially useful PGs are
applied as pharmaceuticals, there emerge the problems that
(1) they are readily transformed metabolically into
inactive substances in living bodies and (2) they exhibit
an unsatisfactory focus selectivity. As the result, the
15 PGs preparations in general have drawbacks in that they
require frequent administration and thus give greater
pain to the patients and moreover their actions to other
tissues than aimed at manifest themselves as side effects.

 The inventors made extensive studies on
20 pharmaceutical application of PGs to overcome the above
difficulties and found previously that a preparation made
by the inclusion of PGE₁ in a fat emulsion for intravenous
administration permits of intravenous administration
accompanied with reduced manifestation of side effects
25 [Japanese Patent Application Kokai (Laid-open) No. 222014/83].

1 On further study the inventors have succeeded, by the
inclusion of PGE_1 alkyl ester (hereinafter referred to
briefly as PGE_1E) in the fat emulsion, in developing a
preparation which has a prolonged half-life of PGs in
5 living body as well as a satisfactory focus selectivity,
and thus accomplished this invention.


PGE_1E is more liposoluble than PGE_1 and hence
a larger amount of it can be incorporated into the fat
emulsion, so that a higher activity can be expected of
10 its fat emulsion even in a smaller dose than that of a fat
emulsion containing PGE_1 .

An object of this invention is to provide a
 PGE_1E fat emulsion for intravenous administration which
releases its effective ingredient sustainedly and at the
15 same time has a good focus selectivity.

Other objects and advantages of this invention
will become apparent from the following description.

The accompanying drawing shows the degree of
decrease in blood pressure observed when the preparation
20 of this invention or a control preparation is separately
administered intravenously. In the drawing, line B
indicates the degree of decrease in blood pressure of the
preparation of this invention and line A indicates that
of the control preparation.

25 In this invention, PGE_1E refers to a compound
represented by the general formula



wherein R denotes an alkyl group having 1 to 30 carbon atoms.

The alkyl group in the above general formula may be of either straight chain or branched chain. The number of its carbon atoms is 1 to 30, preferably 1 to 15 and more preferably 3 to 10. Examples of such alkyl groups include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, n-nonyl and n-decyl.

The PGE_1 content of the present fat emulsion can be suitably varied according to the composition and use of the emulsion, but it should cover the effective amount which is in the range of 100 to 0.2 $\mu\text{g/ml}$.

The fat emulsion, as herein referred to, may comprise, as main constituents, 5 - 50% (W/V) of soybean oil, 1 - 50, preferably 5 - 30, parts by weight of a phospholipid for 100 parts by weight of the soybean oil, and a proper quantity of water. In addition, the fat emulsion may contain, if necessary, emulsifying adjuvant [for example, 0.01 - 0.3% (W/V) of a fatty acid having 6 - 22, preferably 12 - 20, carbon atoms or a physiologically acceptable salt thereof], stabilizers [for example, 0.001 - 0.5, preferably 0.005 - 0.1, % (W/V) of a cholesterol or 0.01 - 5, preferably 0.05 - 1, % (W/V) of a phosphatidic acid], high-molecular-

1. weight stabilizing adjuvant [for example, 0.1 - 5, preferably 0.5 - 1, parts by weight of albumin, dextran, vinyl polymers, nonionic surface active agents, gelatin, or hydroxyethylstarch for 1 part by weight of PGE₁E], or
- 5 isotonizing agents (for example, glycerol or glucose in an amount required for the isotonization).

The soybean oil for use in the present emulsion is a highly purified soybean oil, preferably that one (purity: 99.9% or above in terms of total glyceride including tri-, di-, and mono-glyceride) obtained by further purifying common refined soybean oil by steam distillation.

The phospholipid, as herein referred to, is a purified phospholipid such as egg yolk phospholipid or soybean phospholipid, which is obtained by the common fractionation technique using an organic solvent. For instance, it is prepared by slowly adding, with stirring, acetone to a crude yolk phospholipid dissolved in a cold n-hexane-acetone mixture, collecting the insolubles by filtration, repeating the procedure of dissolution followed by precipitation, and finally removing the solvent by distillation. The product comprises phosphatidylcholine and phosphatidylethanolamine as major constituents and minor amounts of other phospholipids such as phosphatidyl-inositol, phosphatidylserine, and sphingomyelin. Various phospholipids can be used each alone or in combinations.

The fatty acids of 6 - 22 carbon atoms for use as emulsifying adjuvant are those suitable for use in pharmaceuticals. They may be of either straight chain or

- 1 branched chain. Most preferred are straight chain fatty acids such as stearic, oleic, linolic, palmitic, linolenic, and myristic acids. The salts should be physiologically acceptable ones such as, for example, salts with alkali
- 5 metals such as sodium and potassium or with alkaline earth metals such as calcium.

The cholesterol and the phosphatidic acid for use as a stabilizer are those which are suitable for use in pharmaceuticals.

- 10 Suitable high-molecular-weight substances for use as stabilizing adjuvant are as follows: The albumin should be of the human origin, in view of the problem of antigenicity. Suitable vinyl polymers include poly-vinylpyrrolidone.

- 15 Suitable nonionic surface active agents are poly-alkylene glycols (for example, polyethylene glycol having an average molecular weight of 1,000 - 10,000, preferably 4,000 - 6,000), polyoxyalkylene copolymers (for example, a polyoxyethylene-polyoxypropylene copolymer
- 20 having an average molecular weight of 1,000 - 20,000, preferably 6,000 - 10,000), polyoxyalkylene derivatives of hardened castor oil [for example, hardened castor oil polyoxyethylene-(40), or -(20), or -(100) ether], and polyoxyalkylene derivatives of castor oil [for example,
- 25 castor oil polyoxyethylene-(20), or -(40), or -(100) ether].

The present fat emulsion is produced, for example, in the following manner: Predetermined amounts of PGE_1E , phospholipid, and, if necessary, the aforementioned

0132027

1 additives are mixed with soybean oil and the mixture is
heated at 40° to 75°C to accelerate dissolution, whereby
a homogeneous solution is formed. The solution is mixed
with a necessary quantity of water and emulsified at 20°
5 to 80°C by means of a common mixer (e.g. a homomixer) to
form a coarse emulsion. A stabilizer and an isotonizing
agent may be added at this stage. The coarse emulsion is
then subjected to size diminution treatment at 20° to 80°C
by using a homogenizer (e.g. a homogenizer of the high
10 pressure-jet type such as Manton-Gaulin homogenizer or of
the ultrasonic type), resulting in a homogenized, finely
dispersed fat emulsion containing PGE₁E. This emulsion has
an excellent storage stability and the average particle
size is 1.0 μ or below. The homogenization of a coarse
15 emulsion by means of Manton-Gaulin homogenizer is carried
out by passing the coarse emulsion 1 to 2 times through
the homogenizer under a first-stage pressure of 100 - 150
kg/cm² and then 5 to 15 times under a second-stage pressure
of 400 - 700 kg/cm².

20 The present fat emulsion is suitable for administration
through a parenteral route, preferably intravenously. For
instance, a dose of 1 to 100 μ g in terms of PGE₁ is ad-
ministered once a day by the continuous intravenous infusion
at a rate of 0.02 - 0.2 ng/kg body weight per minute.

25 Since the fat emulsion of this invention has a
strong medicinal action, focus selectivity, and is of
sustained release it permits effective treatment of the
patient with a small dose.

1 Further, the present emulsion does not undergo
inactivation which is liable to occur with conventional PG
preparations such as an α -cyclodextrin clathrate compound
of PG. As a consequence, it has become possible to ad-
5 ministrated the present emulsion by intravenous injection
which was believed to be impossible with conventional PG
preparations. The present emulsion exhibits a steady
medicinal effect with a small dose, resulting in reduced
side effects. In addition, there is observed none of those
10 swelling, dull pain, redness, and fever which are apt to
occur in the region where a conventional PG preparation was
introduced.

This invention is illustrated below in detail
with reference to Test Examples and Examples of the fat
15 emulsion of this invention, but the invention is not limited
thereto.

Test Example 1

A group of 4 - 6 male adult mongrel dogs each
weighing about 10 kg was used in each test. The dog was
20 anesthetized with sodium pentobarbital (35 mg/kg, intravenous
injection). Sixty minutes after the anesthesia, the blood
pressure (mmHg) was measured. After additional 30 minutes,
the present fat emulsion prepared as in Example 2 described
hereinafter or a control preparation prepared by dissolving
25 PGE₁ methyl ester in physiological saline was administered
intravenously in a dose of 0.1, 0.3 and 1 μ g/kg in terms of
PGE₁ to respective dog groups, and examined for its effect on

1 the blood pressure of the dogs.

The results were as shown in Fig. 1. As is apparent from Fig. 1, the hypotensive action of the preparation of this invention is distinctly stronger than that of the
5 control preparation.

Test Example 2

The LD₅₀ value in intravenous administration of the present preparation prepared as in Example 2 described hereinafter was 200 ml or more/kg body weight for 10%
10 fat emulsion and 150 ml or more/kg body weight for 20% fat emulsion. No hemolyzation was observed at all when the intravenous drip was conducted at a normal rate.

Example 1

To 30 g of purified soybean oil, were added
15 3.6 g of yolk phospholipid, 900 µg of PGE₁ propyl ester, 0.15 g of sodium palmitate, and 0.15 g of phosphatidic acid. The mixture was heated at 45° to 65°C to form a solution. To the solution, was added 200 ml of distilled water, followed by 7.5 g of glycerol of the official grade (Pharmaco-
20 copoeia of Japan). The mixture was made up to 300 ml with distilled water for injection at 20° - 40°C, and coarsely emulsified in "Homomixer". The coarse emulsion was homogenized by passing 10 times through a Manton-Gaulin-type homogenizer under a first-stage pressure of 120 kg/cm² and a
25 total pressure of 500 kg/cm². There was obtained a homogenized, finely dispersed fat emulsion containing PGE₁

- 1 propyl ester. The emulsion, 0.2 - 0.4 μ in average size of dispersed droplets, contained none of the droplets of 1 μ or above in size.

Example 2

- 5 To 35 g of purified soybean oil, were added 3.0 g of soybean phospholipid, 850 μ g of PGE₁ methyl ester, 0.10 g of sodium linolate and 0.15 g of phosphatidic acid. The mixture was heated at 40° to 60°C to form a solution. To the solution, was added 200 ml of distilled water,
10 followed by 7.5 g of glycerol of the official grade (Pharmacopoeia of Japan). The mixture was made up to 300 ml with distilled water for injection at 20° to 40°C, and coarsely emulsified in "Homomixer".

- The coarse emulsion was homogenized by passing
15 10 times through a Manton-Gaulin-type homogenizer under a first-stage pressure of 120 kg/cm² and a total pressure of 500 kg/cm². There was obtained a homogenized, finely dispersed fat emulsion containing PGE₁ methyl ester. The emulsion, 0.2 to 0.4 μ in average size of dispersed drop-
20 lets, contained none of the droplets of 1 μ or above in size.

Example 3


- To 25 g of purified soybean oil, were added 4.0 g of yolk phospholipid, 800 μ g of PGE₁ ethyl ester, 0.20 g
25 of sodium stearate and 0.20 g of cholesterol. The mixture was heated at 50° to 65°C to form a solution. To the

1 solution, was added 200 ml of distilled water, followed by 7.5 g of glycerol of the official grade (Pharmacopoeia of Japan). The mixture was made up to 300 ml with distilled water for injection at 20° to 40°C, and coarsely emulsified in "Homomixer". The coarse emulsion was homogenized by passing 10 times through a Manton-Gaulin-type homogenizer under a first-stage pressure of 120 kg/cm² and a total pressure of 500 kg/cm².

There was obtained a homogenized, finely dispersed 10 fat emulsion containing PGE₁ ethyl ester. The emulsion, 0.2 to 0.4 μ in average size of dispersed droplets, contained none of the droplets of 1 μ or above in size.

CLAIMS:-

1. A fat emulsion containing prostaglandin E₁ alkyl ester represented by the general formula



5

wherein R denotes an alkyl group having 1 to 30 carbon atoms.

2. A fat emulsion according to Claim 1, wherein the alkyl group in the above general formula is methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, n-nonyl, or n-decyl.

3. A fat emulsion according to Claim 1 or 2, which comprises 5 - 50% (W/V) of soybean oil containing an effective amount of prostaglandin E₁ alkyl ester, 1 - 50 parts by weight of a phospholipid for 100 parts by weight of the soybean oil, and water.

4. A fat emulsion according to Claim 1, 2 or 3, which contains as emulsifying adjuvant 0.01 - 0.3% (W/V) of a fatty acid having 6 - 22 carbon atoms or a physiologically acceptable salt thereof.


5. A fat emulsion according to Claim 1, 2, 3 or 4, which contains as stabilizer 0.001 - 0.5% (W/V) of a cholesterol or 0.01 - 5% (W/V) of a phosphatidic acid.

6. A fat emulsion according to any preceding Claim, which contains as stabilizing adjuvant 0.1 - 5 parts by weight of

at least one high-molecular-weight substance selected from the group consisting of albumin, dextran, vinyl polymers, nonionic surface active agents, gelatin, and hydroxyethyl-starch for 1 part by weight of prostaglandin E₁ alkyl ester.

7. A fat emulsion according to any preceding Claim, which contains an isotonizing agent.

8. A method for producing a fat emulsion containing prostaglandin E₁ alkyl ester, which comprises dissolving prostaglandin E₁ alkyl ester represented by the general formula



15


wherein R denotes an alkyl group having 1 to 30 carbon atoms, and a phospholipid in soybean oil, mixing the resulting solution with water to form a coarse emulsion, and homogenizing the coarse emulsion.

9. A method according to Claim 8, wherein a fat emulsion comprising an effective amount of prostaglandin E₁ alkyl ester, 5 - 50% (W/V) of soybean oil, 1 - 50 parts by weight of a phospholipid for 100 parts by weight of the soybean oil, and water is produced.

10. A method according to Claim 8 or Claim 9, wherein the homogenization is performed by passing the coarse emulsion

through a high pressure-jet type homogenizer 1 - 2 times under a first-stage pressure of 100 - 150 kg/cm² and then 5 - 15 times under a second-stage pressure of 400 - 700 kg/cm².

- 5 11. A method according to Claim 8, 9 or 10, wherein the dissolution, mixing, and homogenization are carried out at 20 - 80°C.
12. A method according to Claim 8, 9, 10 or 11, wherein 0.01 - 0.3%(W/V) of a fatty acid having 6 - 22 carbon atoms or
10 a physiologically acceptable salt thereof is added as emulsifying adjuvant in the dissolution step.
13. A method according to ^{any one of to 12} Claims 8/, wherein 0.001 - 0.5%(W/V) of a cholesterol or 0.01 - 5%(W/V) of a phosphatidic acid is added as stabilizer of fat emulsion in
15 the dissolution step.
14. A method according to ^{any one of to 13} Claims 8/, wherein 0.1 - 5 parts by weight of at least one high-molecular-weight substance selected from the group consisting of albumin, dextran, vinyl polymers, nonionic surface active agents,
20 gelatin, and hydroxyethylstarch for 1 part by weight of the prostaglandin E₁ alkyl ester is added in the mixing step.
15. A method according to any one of Claims 8 to 14, wherein an isotonizing agent is added in the mixing step.





European Patent
Office

EUROPEAN SEARCH REPORT

0132027
Application number

EP 84 30 3304

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 3)
X,Y	EP-A-0 097 481 (TAISHO PHARMACEUTICAL CO., LTD.) * Examples; claims *	1-15	A 61 K 9/10 A 61 K 47/00 A 61 K 31/557
X,Y	--- US-A-4 190 669 (J.J. VOORHEES) * Column 2, lines 47-54; claims *	1-15	
Y	--- CHEMICAL ABSTRACTS, vol. 83, no. 20, 17th November 1975, page 277, no. 168488c, Columbus, Ohio, USA; & JP - A - 75 105 815 (YAMANOUCHI PHARMACEUTICAL CO., LTD.) 20-08-1975 * Abstract *	1-15	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 3)
			A 61 K 9/00 A 61 K 47/00 A 61 K 31/00
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 02-10-1984	Examiner BERTE M. J.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.